Is "indications" a technical term? Is it as clear as 'Uses''?

Allergy Tablets

due to hay fever and other upper respiratory allergies. DIRECTIONS: Adults and children 12 years and over--1 tablet every 4 to 6 hours, not to exceed 6 tablets in 24 hours or as directed by a physician. Children 6 to 11 years—one half the dult dose (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours. For children under

6 years, consult a physician. **EACH TABLET CONTAINS:** Chorpheniramine Maleate 4 mg. May also contain (may differ from brand) D&C Yellow No. 10, Lactose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch.

WARNINGS: May cause excitability especially in children. Do not take this product unless directed by a physician, if you have a breathing problem such as emphysema or chronic bronchitis, or if you have glaucoma or difficulty in urination due to enlargement of the prostate gland. May cause drowsiness; alcohol, sedatives and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages, and

accion, securives and transplaces may introduce the violence and assistance or contact a Poison Control Center immediately.

Store at controlled room temperature 2°-30°C (36°-86°F).

Use by expiration date printed on package. Protect from excessive moisture.

For better identification keep tablets in carton until used.

Made in U.S.A.

'Each Tablet Contains' does not provide the same specific information as "Active Ingredient."

Notice how much clearer the warning are in the new label.

Critical

bold.

information that

may be needed

emergency is in

quickly in an

The borders make this label easier to navigate

Drug Facts

Active ingredient (in each tablet)

Purpose

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ sneezing ■ runny nose ■ itchy, watery eyes ■ itchy throat

- Warnings
 Ask a doctor before use if you have
 glaucoma a breathing problem such as emphysema or chronic bronchitis
 trouble urinating due to an enlarged prostate gland

 The pharmacist before use if you are taking tranquilizers or sedat
- Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives

- ASK a Octor of prarmaces before use in you are saving using When using this product

 If drowsiness may occur avoid alcoholic drinks
 alcohol, sedatives, and tranquillares may increase drowsiness
 be careful when driving a motor vehicle or operating machinery
 excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years and over take 2 tablets every 4 to 6 hours not more than 12 tablets in 24 hours children 6 years to under 12 years take 1 tablet every 4 to 6 hours; not more than 6 tablets in 24 hours children under 6 years ask a doctor

Drug Facts (continued)

Other information ■ store at 20-25°C (68-77°F) ■ protect from excessive moisture

Inactive ingredients D&C yellow no. 10, lactose, magnesium stearate, microcrystall

When paragraphs are changed to lists, each item on the list becomes a direct object for the verb in the introductory clause. Unnecessary words are deleted.

Notice that "effective," in the original "Indications," has been deleted in the revised "Uses." Why would the FDA want that word removed?

The revised "Directions" chart is much easier to use than the "Directions" paragraph in the original.

'Inactive ingredients" are now clearly labeled.